

The construction and deformation of Hom-Novikov superalgebras

Bing Sun, Liangyun Chen, Yan Liu

School of Mathematics and Statistics, Northeast Normal University,
Changchun, 130024, CHINA

Abstract

We study a twisted generalization of Novikov superalgebras, called Hom-Novikov superalgebras. It is shown that two classes of Hom-Novikov superalgebras can be constructed from Hom-supercommutative algebras together with derivations and Hom-Novikov superalgebras with Rota-Baxter operators, respectively. We show that quadratic Hom-Novikov superalgebras are Hom-associative superalgebras and the sub-adjacent Hom-Lie superalgebras of Hom-Novikov superalgebras are 2-step nilpotent. Moreover, we develop the 1-parameter formal deformation theory of Hom-Novikov superalgebras.

Key words: Hom-Novikov superalgebras, Hom-Lie superalgebras, Quadratic, Deformations

MSC(2010): 17B30, 17B35, 15A63

1 Introduction

Novikov algebras were introduced in connection with Hamiltonian operators in the formal variational calculus and the Poisson brackets of hydrodynamic-type. They were used to construct the Virasoro-type Lie algebras. When Gel'fand and Dorfman^[7–9] studied the following operator:

$$H_{ij} = \sum_k c_{ijk} u_k^{(1)} + d_{ijk} u_k^{(0)} \frac{d}{dx}, \quad c_{ijk} \in \mathbf{C}, \quad d_{ijk} = c_{ijk} + c_{jik},$$

they gave the definition of Novikov algebras. In 1987, Zel'manov^[29] began to study Novikov algebras and proved that the dimension of finite-dimensional simple Novikov

Corresponding author(L. Chen): chenly640@nenu.edu.cn.

Supported by NNSF of China (Nos. 11171055 and 11471090), NSF of Jilin province (No. 201115006).

algebras over a field of characteristic zero is one. In algebras, what are paid attention to by mathematician are classifications and structures, but so far we haven't got the systematic theory for general Novikov algebras. In 1992, Osborn^[20,21] had finished the classification of infinite simple Novikov algebras with nilpotent elements over a field of characteristic zero and finite simple Novikov algebras with nilpotent elements over a field of characteristic $p > 0$. In 1995, Xu^[22-25] developed his theory and got the classification of simple Novikov algebras over an algebraically closed field of characteristic zero. Bai and Meng^[2-4] did a series of researches on low dimensional Novikov algebras, such as the structure and classification. Chen construct two kinds of Novikov algebras^[5,6]. Recently, people obtained some properties in Novikov superalgebras^[14,19,30].

To be more precise, recall that a left-symmetric superalgebra is a $\mathbb{Z}_2$ -graded vector space A together with a binary operation $\mu : A \times A \rightarrow A$ satisfying

$$(xy)z - x(yz) = (-1)^{|x||y|}((yx)z - y(xz)), \quad (1.1)$$

for $x \in A_{|x|}$, $y \in A_{|y|}$, $z \in A_{|z|}$, $|x|, |y|, |z| \in \mathbb{Z}_2$. Here and in what follows we often write $\mu(x, y) = xy$. In the other words, if $(x, y, z) = (xy)z - x(yz)$ denotes the associator, then (1.1) shows that (x, y, z) is supersymmetric with respect to two left variables x and y , hence named left-symmetric superalgebra. A Novikov superalgebra is a left-symmetric superalgebra A that satisfies the additional property

$$(xy)z = (-1)^{|y||z|}(xz)y, \text{ for } x \in A, y \in A_{|y|}, z \in A_{|z|}. \quad (1.2)$$

A quadratic Novikov superalgebra, introduced in [19], is a Novikov superalgebra with a symmetric nondegenerate even invariant bilinear form. The motivation for studying quadratic Novikov superalgebras comes from the fact that Lie or associative algebras with symmetric nondegenerate even invariant bilinear forms have important applications in several areas of mathematics and physics, such as the structure theory of finite-dimensional semi-simple Lie algebras, the theory of complete integrable Hamiltonian systems and the classification of statistical models over two-dimensional graphs.

Yau in [26] introduced Hom-Novikov algebras, in which the two defining identities are twisted by a linear map. It turned out that Hom-Novikov algebras can be constructed from Novikov algebras, commutative Hom-associative algebras and Hom-Lie algebras along with some suitable linear maps. And Yuan and You in [27] introduced quadratic Hom-Novikov algebras. Later, Zhang, Hou and Bai in [28] defined a Hom-Novikov superalgebra as a twisted generalization of Novikov superalgebras.

The deformation is a tool to study a mathematical object by deforming it into a family of the same kind of objects depending on a certain parameter. The deformation theory was introduced by Gerstenhaber for rings and algebras^[10-13], by Kubo and Taniguchi for Lie triple systems^[15], by Ma, Chen and Lin for Hom-Lie Yamaguti algebras^[17] and Hom-Lie triple systems^[18], by Bai and Meng for Novikov algebras^[4]. They studied 1-parameter formal deformations and established the connection between the cohomology groups and infinitesimal deformations.

The purpose of this paper is to consider the realization and the 1-parameter formal deformation theory of Hom-Novikov superalgebras based on some work in [4, 18, 26, 27]. The paper is organized as follows. Section 2 concerns the realization of Hom-Novikov superalgebras. It is shown that two classes of Hom-Novikov superalgebras can be constructed from Hom supercommutative algebras together with derivations and Hom-Novikov superalgebras with Rota-Baxter operators, respectively. In Section 3, we introduce the notion of quadratic Hom-Novikov superalgebras and the relationship between Hom-Novikov superalgebras and sub-adjacent Hom-Lie superalgebras. We also show quadratic Hom-Novikov superalgebras are Hom-associative superalgebras and the sub-adjacent Hom-Lie superalgebras of Hom-Novikov superalgebras are 2-step nilpotent. Section 4 is dedicated to the 1-parameter formal deformation theory of Hom-Novikov superalgebras. Moreover, we define low orders coboundary operator and give low order cohomology groups of Hom-Novikov superalgebras. We show that the cohomology group is suitable for this 1-parameter formal deformation theory.

Throughout this paper $\mathbf{F}$ denotes an arbitrary field.

2 Hom-Novikov superalgebra

Let $(A, \cdot)$ be an algebra over field $\mathbf{F}$. A is said to be a superalgebra if the underlying vector space of A is $\mathbb{Z}_2$ -graded (i.e., $A = A_0 \oplus A_1$, where A_0 and A_1 are vector subspaces of A) and $A_\alpha \cdot A_\beta \subset A_{\alpha+\beta}$, $\forall \alpha, \beta \in \mathbb{Z}_2$. An element $x \in A$ is called homogeneous if $x \in A_0 \cup A_1$. In this work, all elements are supposed to be homogeneous unless otherwise stated. For a homogeneous element x we shall use the standard notation $|x| \in \mathbb{Z}_2 = \{\bar{0}, \bar{1}\}$ to indicate its degree.

Definition 2.1. ^[1] A multiplicative Hom-Lie superalgebra is a triple $(A, [\cdot, \cdot]_A, \alpha)$ consisting of a $\mathbb{Z}_2$ -graded vector space L , an even bilinear map $[\cdot, \cdot] : A \times A \rightarrow A$ and an even algebraic morphism $\alpha : A \rightarrow A$ satisfying

$$\alpha([x, y]) = [\alpha(x), \alpha(y)], \quad (2.1)$$

$$[x, y] = -(-1)^{|x||y|}[y, x], \quad (2.2)$$

$$(-1)^{|x||z|}[\alpha(x), [y, z]] + (-1)^{|y||x|}[\alpha(y), [z, x]] + (-1)^{|z||y|}[\alpha(z), [x, y]] = 0, \quad (2.3)$$

where x, y and z are homogeneous elements in A . Lie superalgebras are examples of Hom-Lie superalgebras in which α is the identity map.

Definition 2.2. ^[1] A Hom-associative superalgebra consists of a $\mathbb{Z}_2$ -graded vector space A , a linear self-map α and an even bilinear map $\mu : A \times A \rightarrow A$, satisfying

$$\alpha(xy) = \alpha(x)\alpha(y) \quad (\text{multiplicativity})$$

and

$$(xy)\alpha(z) = \alpha(x)(yz) \quad (\text{Hom-associativity}),$$

for $x, y, z \in A$.

Definition 2.3. ^[28] A Hom-Novikov superalgebra is a triple (A, μ, α) consisting of a $\mathbb{Z}_2$ -graded vector space A , an even bilinear map $\mu : A \times A \rightarrow A$ and an even homomorphism $\alpha : A \rightarrow A$ satisfying

$$\alpha(xy) = \alpha(x)\alpha(y), \quad (2.4)$$

$$(xy)\alpha(z) - \alpha(x)(yz) = (-1)^{|x||y|}((yx)\alpha(z) - \alpha(y)(xz)), \quad (2.5)$$

$$(xy)\alpha(z) = (-1)^{|y||z|}(xz)\alpha(y). \quad (2.6)$$

We see that Novikov superalgebras are examples of Hom-Novikov superalgebras in which α is the identity map. For a Hom-Novikov superalgebra (A, μ, α) , we call μ the Hom-Novikov product of A . Comparing with Definition 2.3 with (1.1) and (1.2), if and only if (2.4) and (2.5) are satisfied, then we call (A, μ, α) a Hom-left-symmetric superalgebra. In particular, a Hom-Novikov superalgebra is a Hom-left-symmetric superalgebra that also satisfies (2.6).

Definition 2.4. Let (A, μ, α) be a Hom-Novikov superalgebra, which is called

- (i) regular if α is an algebra automorphism;
- (ii) involutive if α is an involution, i.e., $\alpha^2 = \text{id}$.

Lemma 2.5. ^[28] Let (A, μ, α) be a Hom-Novikov superalgebra and $[\cdot, \cdot] : A \times A \rightarrow A$ be a binary operation on A defined by

$$[x, y] = xy - (-1)^{|x||y|}yx \quad \forall x, y \in A.$$

Then $HLie(A) = (A, [\cdot, \cdot], \alpha)$ is a Hom-Lie superalgebra with the same twisting map α , which is called the sub-adjacent Hom-Lie superalgebra of A .

Zhang, Hou and Bai in [28] gave a way to construct Hom-Novikov superalgebras, starting from a Novikov superalgebra and an algebra endomorphism. In the following, we provide a construction of Novikov superalgebras from Hom-Novikov superalgebras along with an algebra automorphism.

Theorem 2.6. If (A, μ, α) is an involutive Hom-Novikov superalgebra, then $(A, \alpha \circ \mu, \alpha)$ is a Novikov superalgebra.

Proof. For convenience, we write $x * y = \alpha(xy)$, for all $x, y \in A$. Hence, it needs to show

$$(x * y) * z - x * (y * z) = (-1)^{|x||y|}((y * x) * z - y * (x * z)), \quad (2.7)$$

$$(x * y) * z = (-1)^{|y||z|}(x * z) * y,$$

for all $x, y, z \in A$. Since (A, μ, α) is an involutive Hom-Novikov superalgebra, we have

$$(x * y) * z = \alpha(\alpha(xy)z) = \alpha^2(xy)\alpha(z) = (xy)\alpha(z) = (-1)^{|y||z|}(xz)\alpha(y) = (-1)^{|y||z|}(x * z) * y.$$

Furthermore,

$$\begin{aligned}
(x * y) * z - x * (y * z) &= \alpha(\alpha(xy)z) - \alpha(x\alpha(yz)) \\
&= (xy)\alpha(z) - \alpha(x)(yz) \\
&= (-1)^{|x||y|}((yx)\alpha(z) - \alpha(y)(xz)) \\
&= (-1)^{|x||y|}((y * x) * z - y * (x * z)),
\end{aligned}$$

which proves Equation (2.7) and the proposition. $\square$

Theorem 2.7. *If (A, μ, α) is a regular Hom-Novikov superalgebra, then $(A, [\cdot, \cdot]_{\alpha^{-1}} = \alpha^{-1} \circ [\cdot, \cdot])$ is a Lie superalgebra, where $[x, y] = xy - (-1)^{|x||y|}yx$, for $x, y \in A$. In particular, if α is an involution, then $(A, [\cdot, \cdot]_{\alpha} = \alpha \circ [\cdot, \cdot])$ is a Lie superalgebra.*

Proof. For any $x, y, z \in A$, we have

$$\begin{aligned}
&(-1)^{|x||z|}[x, [y, z]_{\alpha^{-1}}]_{\alpha^{-1}} \\
&= (-1)^{|x||z|}\alpha^{-1}(x[y, z]_{\alpha^{-1}} - (-1)^{|x||y|+|x||z|}[y, z]_{\alpha^{-1}}x) \\
&= (-1)^{|x||z|}\alpha^{-1}(x\alpha^{-1}(yz - (-1)^{|y||z|}zy) - (-1)^{|x||y|+|x||z|}\alpha^{-1}(yz - (-1)^{|y||z|}zy)x) \\
&= \alpha^{-2}((-1)^{|x||z|}\alpha(x)(yz) - (-1)^{|x||z|+|y||z|}\alpha(x)(zy) - (-1)^{|x||y|}(yz)\alpha(x) + (-1)^{|x||y|+|y||z|}(zy)\alpha(x)).
\end{aligned}$$

Similarly, we have

$$\begin{aligned}
&(-1)^{|x||y|}[y, [z, x]_{\alpha^{-1}}]_{\alpha^{-1}} \\
&= \alpha^{-2}((-1)^{|x||y|}\alpha(y)(zx) - (-1)^{|x||y|+|x||z|}\alpha(y)(xz) - (-1)^{|y||z|}(zx)\alpha(y) + (-1)^{|y||z|+|x||z|}(xz)\alpha(y)), \\
&(-1)^{|y||z|}[z, [x, y]_{\alpha^{-1}}]_{\alpha^{-1}} \\
&= \alpha^{-2}((-1)^{|y||z|}\alpha(z)(xy) - (-1)^{|x||y|+|y||z|}\alpha(z)(yx) - (-1)^{|x||z|}(xy)\alpha(z) + (-1)^{|x||y|+|x||z|}(yx)\alpha(z)).
\end{aligned}$$

Then it follows from Equation (2.5) that

$$(-1)^{|x||z|}[x, [y, z]_{\alpha^{-1}}]_{\alpha^{-1}} + (-1)^{|x||y|}[y, [z, x]_{\alpha^{-1}}]_{\alpha^{-1}} + (-1)^{|y||z|}[z, [x, y]_{\alpha^{-1}}]_{\alpha^{-1}} = 0.$$

Clearly, $[x, y] = -(-1)^{|x||y|}[y, x]$. Since α^{-1} is an automorphism, we have $[x, y]_{\alpha^{-1}} = -(-1)^{|x||y|}[y, x]_{\alpha^{-1}}$, which proves that $(A, [\cdot, \cdot]_{\alpha^{-1}})$ is a Lie superalgebra. It follows immediately that $(A, [\cdot, \cdot]_{\alpha})$ is also a Lie superalgebra since $\alpha = \alpha^{-1}$ when α is an involution. $\square$

Theorem 2.8. *Let (A, μ, α) be a Hom-Novikov superalgebra. Then $(A, \alpha \circ \mu, \alpha^2)$ is a Hom-Novikov superalgebra.*

Proof. For convenience, we write $x * y = \alpha(xy)$, for all $x, y \in A$. Hence, it needs to show

$$(x * y) * \alpha^2(z) - \alpha^2(x) * (y * z) = (-1)^{|x||y|}((y * x) * \alpha^2(z) - \alpha^2(y) * (x * z)), \quad (2.8)$$

$$(x * y) * \alpha^2(z) = (-1)^{|y||z|}(x * z) * \alpha^2(y),$$

for all $x, y, z \in A$. Since (A, μ, α) is a Hom-Novikov superalgebra, we have

$$(x * y) * \alpha^2(z) = \alpha^2((xy)\alpha(z)) = (-1)^{|y||z|} \alpha^2((xz)\alpha(y)) = (-1)^{|y||z|} (x * z) * \alpha^2(y).$$

Furthermore,

$$\begin{aligned} (x * y) * \alpha^2(z) - \alpha^2(x) * (y * z) &= \alpha^2((xy)\alpha(z) - \alpha(x)(yz)) \\ &= (-1)^{|x||y|} \alpha^2((yx)\alpha(z) - \alpha(y)(xz)) \\ &= (-1)^{|x||y|} ((y * x) * \alpha^2(z) - \alpha^2(y) * (x * z)), \end{aligned}$$

which proves Equation (2.8) and the proposition. $\square$

We now construct several classes of Hom-Novikov superalgebras. Yau in [26] gave such construction of Hom-Novikov algebras, starts with an Hom-associative and commutative algebra (A, μ) and an even derivation $D : A \rightarrow A$. The new product

$$x * y = \mu(x, D(y)) = xD(y), \quad (2.9)$$

for $x, y \in A$ makes $(A, *)$ into a Hom-Novikov algebra. To generalize this construction, we define a Hom supercommutative algebra to be a Hom-associative superalgebra whose multiplication is supercommutative. An even derivation on a Hom-associative superalgebra is defined in the usual way. Then we have the following result.

Theorem 2.9. *Let (A, μ, α) be a Hom-supercommutative algebra and $D : A \rightarrow A$ be an even derivation such that $D\alpha = \alpha D$. Then $(A, *, \alpha)$ is a Hom-Novikov superalgebra, where $*$ is defined as in (2.9).*

Proof. The multiplication of α with respect to $*$ in (2.9) follows from the multiplication of α with respect to μ and the hypothesis $D\alpha = \alpha D$. Next we check (2.5)

$$\begin{aligned} &(x * y) * \alpha(z) - \alpha(x) * (y * z) \\ &= (xD(y))D(\alpha(z)) - \alpha(x)D(yD(z)) \\ &= (xD(y))\alpha(D(z)) - \alpha(x)(D(y)D(z)) - \alpha(x)(yD^2(z)) \\ &= -\alpha(x)(yD^2(z)) \\ &= -(xy)\alpha(D^2(z)). \end{aligned}$$

The last two equalities follow from Hom-associativity. On the other hand,

$$\begin{aligned} &(-1)^{|x||y|} ((y * x) * \alpha(z) - \alpha(y) * (x * z)) \\ &= (-1)^{|x||y|} ((yD(x))D(\alpha(z)) - \alpha(y)D(xD(z))) \\ &= (-1)^{|x||y|} ((yD(x))\alpha(D(z)) - \alpha(y)(D(x)D(z)) - \alpha(y)(xD^2(z))) \\ &= (-1)^{|x||y|} (\alpha(y)(D(x)D(z)) - \alpha(y)(D(x)D(z)) - \alpha(y)(xD^2(z))) \\ &= -(-1)^{|x||y|} \alpha(y)(xD^2(z)) \\ &= -(-1)^{|x||y|} (yx)\alpha(D^2(z)) \end{aligned}$$

$$= -(xy)\alpha(D^2(z)).$$

Futhermore, we have

$$\begin{aligned} (x * y) * \alpha(z) &= (xD(y))\alpha(D(z)) = \alpha(x)(D(y)D(z)) \\ &= (-1)^{|y||z|}\alpha(x)(D(z)D(y)) \\ &= (-1)^{|y||z|}(xD(z))\alpha(D(y)) \\ &= (-1)^{|y||z|}(x * z) * \alpha(y). \end{aligned}$$

Consequently, we prove the theorem. $\square$

Theorem 2.10. *Let (A, μ) be an associative superalgebra whose multiplication is supercommutative, $\alpha : A \rightarrow A$ be an even algebra morphism, and $D : A \rightarrow A$ be an even derivation such that $D\alpha = \alpha D$. Then $(A, *, \alpha)$ is a Hom-Novikov superalgebra, where*

$$x * y = \alpha(xD(y)), \quad \forall x, y \in A.$$

Proof. First, we have

$$\alpha(x * y) = \alpha \circ \alpha(xD(y)) = \alpha(\alpha(x)D(\alpha(y))) = \alpha(x) * \alpha(y).$$

On one hand, we have

$$\begin{aligned} &(x * y) * \alpha(z) - \alpha(x) * (y * z) \\ &= \alpha(xD(y)) * \alpha(z) - \alpha(x) * \alpha(yD(z)) \\ &= \alpha((xD(y)) * z - x * (yD(z))) \\ &= \alpha(\alpha((xD(y))D(z)) - \alpha(xD(yD(z)))) \\ &= \alpha^2((xD(y))D(z) - xD(yD(z))) \\ &= \alpha^2(x(D(y)D(z)) - x(D(y)D(z)) - x(yD^2(z))) \\ &= -\alpha^2(x(yD^2(z))) \\ &= -\alpha^2((xy)D^2(z)). \end{aligned}$$

Similarly, we have

$$\begin{aligned} &(-1)^{|x||y|}((y * x) * \alpha(z) - \alpha(y) * (x * z)) \\ &= (-1)^{|x||y|}\alpha((yD(x)) * z - y * (xD(z))) \\ &= (-1)^{|x||y|}\alpha^2((yD(x))D(z) - yD(xD(z))) \\ &= (-1)^{|x||y|}\alpha^2(y(D(x)D(z)) - y(D(x)D(z)) - y(xD^2(z))) \\ &= -(-1)^{|x||y|}\alpha^2((yx)D^2(z)) \\ &= -\alpha^2((xy)D^2(z)). \end{aligned}$$

Futhermore, we have

$$(x * y) * \alpha(z) = \alpha(xD(y)) * \alpha(z) = \alpha^2((xD(y))D(z))$$

$$\begin{aligned}
&= \alpha^2(x(D(y)D(z))) \\
&= (-1)^{|y||z|}\alpha^2(x(D(z)D(y))) \\
&= (-1)^{|y||z|}\alpha^2((xD(z)D(y))) \\
&= (-1)^{|y||z|}(x * z) * \alpha(y).
\end{aligned}$$

Consequently, we prove the theorem. $\square$

Definition 2.11. Let $(A, *, \alpha)$ be a Hom-superalgebra and let $\lambda \in \mathbf{F}$. If a linear map $P : A \rightarrow A$ satisfies

$$P(x) * P(y) = P(P(x) * y + x * P(y) + \lambda x * y), \quad \forall x, y \in A,$$

then P is called a Rota-Baxter operator of weight λ and $(A, *, \alpha, P)$ is called a Rota-Baxter Hom-superalgebra of weight λ .

Theorem 2.12. Let $(A, *, \alpha, P)$ be a Rota-Baxter Hom-Novikov superalgebra of weight λ and P an even linear map. Assume that α and P commute. Then $(A, \circ, \alpha, P)$ is a Hom-Novikov superalgebra, where the multiplication $\circ$ is defined as

$$x \circ y := P(x) * y + x * P(y) + \lambda x * y, \quad \forall x, y \in A.$$

Proof. The multiplication of α with respect to $\circ$ follows from the multiplication of α with respect to $*$ and the hypothesis $P\alpha = \alpha P$. For any $x, y, z \in A$, we have,

$$\begin{aligned}
&(x \circ y) \circ \alpha(z) - \alpha(x) \circ (y \circ z) \\
&= P(P(x) * y + x * P(y) + \lambda x * y) * \alpha(z) + (P(x) * y + x * P(y) + \lambda x * y) * P(\alpha(z)) \\
&\quad + \lambda(P(x) * y + x * P(y) + \lambda x * y) * \alpha(z) - P(\alpha(x)) * (P(y) * z + y * P(z) + \lambda y * z) \\
&\quad - \alpha(x) * P(P(y) * z + y * P(z) + \lambda y * z) - \lambda \alpha(x) * (P(y) * z + y * P(z) + \lambda y * z) \\
&= (P(x) * P(y)) * \alpha(z) + (P(x) * y) * \alpha(P(z)) + (x * P(y)) * \alpha(P(z)) + \lambda(x * y) * \alpha(P(z)) \\
&\quad + \lambda(P(x) * y) * \alpha(z) + \lambda(x * P(y)) * \alpha(z) + \lambda^2(x * y) * \alpha(z) - \alpha(P(x)) * (P(y) * z) \\
&\quad - \alpha(P(x)) * (y * P(z)) - \lambda \alpha(P(x)) * (y * z) - \alpha(x) * (P(y) * P(z)) - \lambda \alpha(x) * (P(y) * z) \\
&\quad - \lambda \alpha(x) * (y * P(z)) - \lambda^2 \alpha(x) * (y * z) \\
&= (P(x) * P(y)) * \alpha(z) - \alpha(P(x)) * (P(y) * z) + (P(x) * y) * \alpha(P(z)) - \alpha(P(x)) * (y * P(z)) \\
&\quad + (x * P(y)) * \alpha(P(z)) - \alpha(x) * (P(y) * P(z)) + \lambda(x * y) * \alpha(P(z)) - \lambda \alpha(x) * (y * P(z)) \\
&\quad + \lambda(P(x) * y) * \alpha(z) - \lambda \alpha(P(x)) * (y * z) + \lambda(x * P(y)) * \alpha(z) - \lambda \alpha(x) * (P(y) * z) \\
&\quad + \lambda^2(x * y) * \alpha(z) - \lambda^2 \alpha(x) * (y * z) \\
&= (-1)^{|x||y|} \left((P(y) * P(x)) * \alpha(z) - \alpha(P(y)) * (P(x) * z) + (y * P(x)) * \alpha(P(z)) \right. \\
&\quad \left. - \alpha(y) * (P(x) * P(z)) + (P(y) * x) * \alpha(P(z)) - \alpha(P(y)) * (x * P(z)) + \lambda(y * x) * \alpha(P(z)) \right. \\
&\quad \left. - \lambda \alpha(y) * (x * P(z)) + \lambda(y * P(x)) * \alpha(z) - \lambda \alpha(y) * (P(x) * z) + \lambda(P(y) * x) * \alpha(z) \right. \\
&\quad \left. - \lambda \alpha(P(y)) * (x * z) + \lambda^2(y * x) * \alpha(z) - \lambda^2 \alpha(y) * (x * z) \right).
\end{aligned}$$

Similarly, we have

$$\begin{aligned}
& (-1)^{|x||y|} \left((y \circ x) \circ \alpha(z) - \alpha(y) \circ (x \circ z) \right) \\
&= (-1)^{|x||y|} \left((P(y) * P(x)) * \alpha(z) - \alpha(P(y)) * (P(x) * z) + (P(y) * x) * \alpha(P(z)) \right. \\
&\quad - \alpha(P(y)) * (x * P(z)) + (y * P(x)) * \alpha(P(z)) - \alpha(y) * (P(x) * P(z)) + \lambda(y * x) * \alpha(P(z)) \\
&\quad - \lambda\alpha(y) * (x * P(z)) + \lambda(P(y) * x) * \alpha(z) - \lambda\alpha(P(y)) * (x * z) + \lambda(y * P(x)) * \alpha(z) \\
&\quad \left. - \lambda\alpha(y) * (P(x) * z) + \lambda^2(y * x) * \alpha(z) - \lambda^2\alpha(y) * (x * z) \right).
\end{aligned}$$

Furthermore, on the one hand, we have

$$\begin{aligned}
& (x \circ y) \circ \alpha(z) \\
&= P(P(x) * y + x * P(y) + \lambda x * y) * \alpha(z) + (P(x) * y + x * P(y) + \lambda x * y) * P(\alpha(z)) \\
&\quad + \lambda(P(x) * y + x * P(y) + \lambda x * y) * \alpha(z) \\
&= (P(x) * P(y)) * \alpha(z) + (P(x) * y) * \alpha(P(z)) + (x * P(y)) * \alpha(P(z)) + \lambda(x * y) * \alpha(P(z)) \\
&\quad + \lambda(P(x) * y) * \alpha(z) + \lambda(x * P(y)) * \alpha(z) + \lambda^2(x * y) * \alpha(z) \\
&= (-1)^{|y||z|} \left((P(x) * z) * \alpha(P(y)) + (P(x) * P(z)) * \alpha(y) + (x * P(z)) * \alpha(P(y)) \right. \\
&\quad \left. + \lambda(x * P(z)) * \alpha(y) + \lambda(P(x) * z) * \alpha(y) + \lambda(x * z) * \alpha(P(y)) + \lambda^2(x * z) * \alpha(y) \right).
\end{aligned}$$

On the other hand, we have

$$\begin{aligned}
& (-1)^{|y||z|} \left((x \circ z) \circ \alpha(y) \right) \\
&= (-1)^{|y||z|} \left((P(x) * P(z)) * \alpha(y) + (P(x) * z) * \alpha(P(y)) + (x * P(z)) * \alpha(P(y)) \right. \\
&\quad \left. + \lambda(x * z) * \alpha(P(y)) + \lambda(P(x) * z) * \alpha(y) + \lambda(x * P(z)) * \alpha(y) + \lambda^2(x * z) * \alpha(y) \right).
\end{aligned}$$

Hence, the conclusion holds. $\square$

3 Quadratic Hom-Novikov superalgebras

Definition 3.1. ^[16] Let A be a Hom-Lie superalgebra. A bilinear form B on A is

- (i) *supersymmetric* if $B(x, y) = (-1)^{|x||y|} B(y, x)$, $\forall x, y \in A$;
- (ii) *nondegenerate* if $x, y \in A$ satisfies $B(x, y) = 0$, $\forall y \in A$, then $x = 0$;
- (iii) *invariant* if $B([x, y], z) = B(x, [y, z])$, $\forall x, y, z \in A$;
- (iv) *even* if $B(A_{\bar{0}}, A_{\bar{1}}) = B(A_{\bar{1}}, A_{\bar{0}}) = \{0\}$.

Definition 3.2. ^[16] A quadratic Hom-Lie superalgebra is a quadruple $(A, [\cdot, \cdot], \alpha, B)$ such that $(A, [\cdot, \cdot], \alpha)$ is a Hom-Lie superalgebra with an even, supersymmetric, non-degenerate and invariant bilinear form B on A satisfying

$$B(\alpha(x), y) = B(x, \alpha(y)), \quad \forall x, y \in A. \quad (3.1)$$

We recover quadratic Lie superalgebras when $\alpha = \text{id}$.

Definition 3.3. A quadratic Hom-Novikov superalgebra is a quadruple (A, μ, α, B) such that (A, μ, α) is a Hom-Novikov superalgebra with an even, supersymmetric, non-degenerate bilinear form B on A satisfying

$$B(\alpha(x), yz) = B(xy, \alpha(z)), \quad \forall x, y, z \in A. \quad (3.2)$$

We recover quadratic Novikov superalgebras when $\alpha = \text{id}$.

Theorem 3.4. Let (A, μ, α, B) be a quadratic Hom-Novikov superalgebra and $HLie(A) = (A, [\cdot, \cdot], \alpha)$ the sub-adjacent Hom-Lie superalgebra of A . If α is an automorphism satisfying

$$B(\alpha(x), y) = B(x, \alpha(y)), \quad \forall x, y \in A, \quad (3.3)$$

then $(A, [\cdot, \cdot], \alpha, B_\alpha)$ is a quadratic Hom-Lie superalgebra, where $B_\alpha(x, y) = B(\alpha(x), y)$.

Proof. Since B is a nondegenerate even bilinear form and α is an automorphism, B_α is a nondegenerate even bilinear form on A . For all $x, y, z \in A$, using the properties of B , we have

$$\begin{aligned} B_\alpha([x, y], z) &= B(\alpha([x, y]), z) = B([x, y], \alpha(z)) \\ &= B(xy, \alpha(z)) - (-1)^{|x||y|} B(yx, \alpha(z)) \\ &= B(\alpha(x), yz) - (-1)^{|x||y|} (-1)^{|x||y|+|y||z|} B(\alpha(x), zy) \\ &= B(\alpha(x), [y, z]) = B_\alpha(x, [y, z]). \end{aligned}$$

Hence B_α is invariant. Using supersymmetry of B and Equation (3.3), we have

$$B_\alpha(x, y) = B(\alpha(x), y) = (-1)^{|x||y|} B(y, \alpha(x)) = (-1)^{|x||y|} B(\alpha(y), x) = (-1)^{|x||y|} B_\alpha(y, x),$$

which proves B_α is supersymmetry. Using Equation (3.3) again, we have

$$B_\alpha(\alpha(x), y) = B(\alpha(\alpha(x)), y) = B(\alpha(x), \alpha(y)) = B_\alpha(x, \alpha(y)),$$

which completes the proof. □

Corollary 3.5. Let (A, μ, B) be a quadratic Novikov superalgebra with an automorphism satisfying Equation (3.3) and $(A, [\cdot, \cdot])$ be the sub-adjacent Lie superalgebra. Then $(A, [\cdot, \cdot]_\alpha = \alpha \circ [\cdot, \cdot], \alpha, B_\alpha)$ forms a quadratic Hom-Lie superalgebra, where $B_\alpha(x, y) = B(\alpha(x), y)$.

Proof. It is obvious that $(A, [\cdot, \cdot]_\alpha, \alpha)$ is a Hom-Lie superalgebra. Using the similar arguments as those in the proof of Theorem 3.4, we get B_α is an even symmetric non-degenerate bilinear form with Equation (3.1) satisfied. It remains to show that B_α is invariant. For all $x, y, z \in A$, using invariance and supersymmetry of B , we have

$$\begin{aligned} B_\alpha([x, y]_\alpha, z) &= B(\alpha([x, y]_\alpha), z) = B([x, y]_\alpha, \alpha(z)) \\ &= B(\alpha(x)\alpha(y), \alpha(z)) - (-1)^{|x||y|} B(\alpha(y)\alpha(x), \alpha(z)) \\ &= B(\alpha(x), \alpha(y)\alpha(z)) - (-1)^{|x||y|} (-1)^{|x||y|+|y||z|} B(\alpha(x), \alpha(z)\alpha(y)) \\ &= B(\alpha(x), [y, z]_\alpha) = B_\alpha(x, [y, z]_\alpha), \end{aligned}$$

which proves the invariance of B_α and the result. $\square$

Theorem 3.6. *Let (A, μ, α, B) be a quadratic Hom-Novikov superalgebra, where α is an involution satisfying Equation (3.3). Then $(A, \alpha \circ \mu, B)$ is a quadratic Novikov superalgebra.*

Proof. $(A, \alpha \circ \mu)$ is a Novikov superalgebra by Theorem 2.6. It suffices to show that B is invariant under the operation $\alpha \circ \mu$. For all $x, y, z \in A$, we have

$$B(x, \alpha(y)\alpha(z)) = B(\alpha(x), yz) = B(xy, \alpha(z)) = B(\alpha(x)\alpha(y), z),$$

which completes the proof. $\square$

Theorem 3.7. *Let (A, μ, α, B) is a quadratic Hom-Novikov superalgebra, where α is an automorphism satisfying Equation (3.3). Then $(A, * = \alpha \circ \mu, \alpha^2, B_{\alpha^2})$ is a quadratic Hom-Novikov superalgebra, where $B_{\alpha^2}(x, y) = B(\alpha^2(x), y)$.*

Proof. It follows from Theorem 2.8 that $(A, *, \alpha^2)$ forms a Hom-Novikov superalgebra. Since B is a nondegenerate even bilinear form on A and α is an automorphism, B_{α^2} is a nondegenerate even bilinear form. For all $x, y, z \in A$, using the hypothesis, we have

$$B_{\alpha^2}(x, y) = B(\alpha^2(x), y) = B(x, \alpha^2(y)) = (-1)^{|x||y|} B(\alpha^2(y), x) = (-1)^{|x||y|} B_{\alpha^2}(y, x),$$

thus, B_{α^2} is supersymmetric. Moreover, we have

$$\begin{aligned} B_{\alpha^2}(\alpha^2(x), y * z) &= B(\alpha^4(x), \alpha(y)\alpha(z)) = B(\alpha^3(x), \alpha^2(y)\alpha^2(z)) = B(\alpha(x)\alpha(y), \alpha^4(z)) \\ &= (-1)^{|x||z|+|y||z|} B(\alpha^4(z), x * y) = B_{\alpha^2}(x * y, \alpha^2(z)), \end{aligned}$$

which proves the invariance of B_{α^2} and the theorem. $\square$

Corollary 3.8. *Let (A, μ, α, B) be a quadratic Hom-Novikov superalgebra, where α is an automorphism satisfying Equation (3.3). Then $(A, \alpha^n \circ \mu, \alpha^n, B_{\alpha^n})$ is a quadratic Hom-Novikov superalgebra, where $B_{\alpha^n}(x, y) = B(\alpha^n(x), y)$, $\forall n > 0$.*

Let (A, μ, α) be a Hom-Novikov superalgebra, whose center is denoted by $Z(A)$ and defined by

$$Z(A) = \{x \in A \mid xy = yx = 0, \forall y \in A\}.$$

Let $(A, [\cdot, \cdot], \beta)$ be a Hom-Lie superalgebra. The lower central series of A is defined as usual, i.e., $A^0 = A$, $A^i = [A, A^{i-1}]$, $\forall i \geq 1$. We call A is i -step nilpotent if $A^i = 0$ and $A^{i-1} \neq 0$. The center of the Hom-Lie superalgebra is denoted by $C(A)$ and defined by

$$C(A) = \{x \in A \mid [x, y] = 0, \forall y \in A\}.$$

Theorem 3.9. *Let (A, μ, α, B) be a quadratic Hom-Novikov superalgebra. If α is an automorphism, then (A, μ, α) is a Hom-associative superalgebra.*

Proof. Define $(x, y, z) = \alpha(x)yz - (xy)\alpha(z)$. For any $x, y, z, d \in A$, we have

$$\begin{aligned} B((x, y, z), \alpha(d)) &= B(\alpha(x)(yz), \alpha(d)) - B((xy)\alpha(z), \alpha(d)) \\ &= B(\alpha^2(x), (yz)d) - B(\alpha(x)\alpha(y), \alpha(z\alpha^{-1}(d))) \\ &= B(\alpha^2(x), (yz)d) - B(\alpha^2(x), \alpha(y)(z\alpha^{-1}(d))) \\ &= -B(\alpha^2(x), (y, z, \alpha^{-1}(d))). \end{aligned}$$

Thus, we have

$$\begin{aligned} B((x, y, z), \alpha(d)) &= -B(\alpha^2(x), (y, z, \alpha^{-1}(d))) \\ &= (-1)^{1+|x|(|y|+|z|+|d|)+|y||z|} B((z, y, \alpha^{-1}(d)), \alpha^2(x)) \\ &= (-1)^{|x|(|y|+|z|+|d|)+|y||z|} B(\alpha^2(z), (y, \alpha^{-1}(d), x)) \\ &= (-1)^{|x|(|y|+|z|+|d|)+|y||z|+|z|(|y|+|d|+|x|)} B((y, \alpha^{-1}(d), x), \alpha^2(z)) \\ &= (-1)^{|x|(|y|+|z|+|d|)+|y||z|+|z|(|y|+|d|+|x|)+|y||d|} B((\alpha^{-1}(d), y, x), \alpha^2(z)) \\ &= (-1)^{1+|x|(|y|+|z|+|d|)+|y||z|+|z|(|y|+|d|+|x|)+|y||d|} B(\alpha(d), (y, x, z)) \\ &= (-1)^{1+|x|(|y|+|z|+|d|)+|y||z|+|z|(|y|+|d|+|x|)+|y||d|+|d|(|x|+|y|+|z|)+|x||y|} B((x, y, z), \alpha(d)) \\ &= -B((x, y, z), \alpha(d)). \end{aligned}$$

It follows that $(x, y, z) = 0$ by the non-degeneracy of B . □

Theorem 3.10. *Let (A, μ, α, B) be a quadratic Hom-Novikov superalgebra and $HLie(A)$ be the sub-adjacent Hom-Lie superalgebra. If α is an automorphism, then $[x, y] \subseteq Z(A)$, for any $x, y \in HLie(A)$. As a consequence, $HLie(A)$ is 2-step nilpotent.*

Proof. For any $x, y, z \in A$, by Theorem 3.9 we have

$$\begin{aligned} \alpha(z)[x, y] &= \alpha(z)(xy) - (-1)^{|x||y|}\alpha(z)(yx) = (zx)\alpha(y) - (-1)^{|x||y|}(zy)\alpha(x) \\ &= (zx)\alpha(y) - (zx)\alpha(y) = 0. \end{aligned}$$

Using Equation (3.2), we have

$$B([x, y]z, \alpha(d)) = B(\alpha[x, y], zd) = (-1)^{(|z|+|d|)(|x|+|y|)} B(zd, \alpha([x, y]))$$

$$=(-1)^{(|z|+|d|)(|x|+|y|)}B(\alpha(z), d[x, y]) = 0,$$

which implies $[x, y] \in Z(A)$ since α is an automorphism and B is nondegenerate. Hence, we have $[HLie(A), HLie(A)] \subseteq Z(A)$. Obviously, $Z(A) \subseteq C(HLie(A))$. Then it follows that $HLie(A)$ is 2-step nilpotent. $\square$

Remark 3.11. *A 2-step nilpotent quadratic Hom-Lie superalgebra A admits a quadratic Hom-Novikov superalgebra. It suffices to define a bilinear product on A by $xy = \frac{1}{2}[x, y]$.*

4 1-parameter formal deformations of Hom-Novikov superalgebras

Definition 4.1. *Let $(A, *, \alpha)$ is a Hom-Novikov superalgebra. If an n -linear map $f : \underbrace{A \times \cdots \times A}_{n \text{ times}} \rightarrow A$ satisfies*

$$\alpha(f(x_1, \cdots, x_n)) = f(\alpha(x_1), \cdots, \alpha(x_n)), \quad (4.1)$$

then f is called an n -Hom-cochain on A . Denote by $C_\alpha^n(A, A)$ the set of all n -Hom-cochains, $\forall n \geq 1$.

Definition 4.2. *For $n = 1, 2$, the coboundary operator $\delta_{hom}^n : C_\alpha^n(A, A) \rightarrow C_\alpha^{n+1}(A, A)$ is defined as follows.*

$$\begin{aligned} \delta_{hom}^1 f(x_1, x_2) &= (-1)^{x_1 f} x_1 * f(x_2) + f(x_1) * x_2 - f(x_1 * x_2); \\ \delta_{hom}^2 f(x_1, x_2, x_3) &= f(\alpha(x_1), x_2 * x_3) - (-1)^{|x_1||x_2|} f(\alpha(x_2), x_1 * x_3) \\ &\quad + (-1)^{|x_1||x_2|} f(x_2 * x_1, \alpha(x_3)) + (-1)^{|x_1||f|} \alpha(x_1) * f(x_2, x_3) \\ &\quad - (-1)^{|x_1||x_2|+|x_2||f|} \alpha(x_2) * f(x_1, x_3) + (-1)^{|x_1||x_2|} f(x_2, x_1) * \alpha(x_3) \\ &\quad - (-1)^{|x_2||x_3|} f(x_1 * x_3, \alpha(x_2)) - (-1)^{|x_2||x_3|} f(x_1, x_3) * \alpha(x_2). \end{aligned}$$

It is not difficult to verify that $\delta_{hom}^1 f, \delta_{hom}^2 f$ satisfies (4.1). Thus, the coboundary operator δ_{hom}^n is well-defined.

Theorem 4.3. *The coboundary operator $\delta_{hom}^1, \delta_{hom}^2$ defined above satisfies $\delta_{hom}^2 \delta_{hom}^1 = 0$.*

Proof. Suppose that $f \in C_\alpha^1(A, A)$, we have

$$\begin{aligned} &\delta_{hom}^2 \delta_{hom}^1 f(x_1, x_2, x_3) \\ &= \delta_{hom}^1 f(\alpha(x_1), x_2 * x_3) - (-1)^{|x_1||x_2|} \delta_{hom}^1 f(\alpha(x_2), x_1 * x_3) \\ &\quad + (-1)^{|x_1||x_2|} \delta_{hom}^1 f(x_2 * x_1, \alpha(x_3)) + (-1)^{|x_1||f|} \alpha(x_1) * \delta_{hom}^1 f(x_2, x_3) \\ &\quad - (-1)^{|x_1||x_2|+|x_2||f|} \alpha(x_2) * \delta_{hom}^1 f(x_1, x_3) + (-1)^{|x_1||x_2|} \delta_{hom}^1 f(x_2, x_1) * \alpha(x_3) \\ &\quad - (-1)^{|x_2||x_3|} \delta_{hom}^1 f(x_1 * x_3, \alpha(x_2)) - (-1)^{|x_2||x_3|} \delta_{hom}^1 f(x_1, x_3) * \alpha(x_2) \end{aligned}$$

$$\begin{aligned}
&= (-1)^{|x_1||f|} \alpha(x_1) * f(x_2 * x_3) + f(\alpha(x_1)) * (x_2 * x_3) - f(\alpha(x_1) * (x_2 * x_3)) \\
&\quad - (-1)^{|x_1||x_2|+|x_2||f|} \alpha(x_2) * f(x_1 * x_3) - (-1)^{|x_1||x_2|} f(\alpha(x_2)) * (x_1 * x_3) \\
&\quad + (-1)^{|x_1||x_2|} f(\alpha(x_2) * (x_1 * x_3)) + (-1)^{|x_1||x_2|+(|x_1|+|x_2|)|f|} (x_2 * x_1) * f(\alpha(x_3)) \\
&\quad + (-1)^{|x_1||x_2|} f(x_2 * x_1) * \alpha(x_3) - (-1)^{|x_1||x_2|} f((x_2 * x_1) * \alpha(x_3)) \\
&\quad + (-1)^{(|x_1|+|x_2|)|f|} \alpha(x_1) * (x_2 * f(x_3)) + (-1)^{|x_1||f|} \alpha(x_1) * (f(x_2) * x_3) \\
&\quad - (-1)^{|x_1||f|} \alpha(x_1) * f(x_2 * x_3) - (-1)^{|x_1||x_2|+(|x_1|+|x_2|)|f|} \alpha(x_2) * (x_1 * f(x_3)) \\
&\quad - (-1)^{|x_1||x_2|+|x_2||f|} \alpha(x_2) * (f(x_1) * x_3) + (-1)^{|x_1||x_2|+|x_2||f|} \alpha(x_2) * f(x_1 * x_3) \\
&\quad + (-1)^{|x_1||x_2|+|x_2||f|} (x_2 * f(x_1)) * \alpha(x_3) + (-1)^{|x_1||x_2|} (f(x_2) * x_1) * \alpha(x_3) \\
&\quad - (-1)^{|x_1||x_2|} f(x_2 * x_1) * \alpha(x_3) - (-1)^{|x_2||x_3|+(|x_1|+|x_3|)|f|} (x_1 * x_3) * f(\alpha(x_2)) \\
&\quad - (-1)^{|x_2||x_3|} f(x_1 * x_3) * \alpha(x_2) + (-1)^{|x_2||x_3|} f((x_1 * x_3) * \alpha(x_2)) \\
&\quad - (-1)^{|x_2||x_3|+|x_1||f|} (x_1 * f(x_3)) * \alpha(x_2) - (-1)^{|x_2||x_3|} (f(x_1) * x_3) * \alpha(x_2) \\
&\quad + (-1)^{|x_2||x_3|} f(x_1 * x_3) * \alpha(x_2) \\
&= f(\alpha(x_1)) * (x_2 * x_3) - (-1)^{|x_1||x_2|+|x_2||f|} \alpha(x_2) * (f(x_1) * x_3) \tag{a1} \\
&\quad + (-1)^{|x_1||x_2|+|x_2||f|} (x_2 * f(x_1)) * \alpha(x_3) \tag{b1} \\
&\quad - f(\alpha(x_1) * (x_2 * x_3)) + (-1)^{|x_1||x_2|} f(\alpha(x_2) * (x_1 * x_3)) \tag{a2} \\
&\quad - (-1)^{|x_1||x_2|} f((x_2 * x_1) * \alpha(x_3)) \tag{b2} \\
&\quad - (-1)^{|x_1||x_2|} f(\alpha(x_2)) * (x_1 * x_3) + (-1)^{|x_1||f|} \alpha(x_1) * (f(x_2) * x_3) \tag{a3} \\
&\quad + (-1)^{|x_1||x_2|} (f(x_2) * x_1) * \alpha(x_3) \tag{b3} \\
&\quad + (-1)^{|x_1||x_2|+(|x_1|+|x_2|)|f|} (x_2 * x_1) * f(\alpha(x_3)) + (-1)^{(|x_1|+|x_2|)|f|} \alpha(x_1) * (x_2 * f(x_3)) \tag{a4} \\
&\quad - (-1)^{|x_1||x_2|+(|x_1|+|x_2|)|f|} \alpha(x_2) * (x_1 * f(x_3)) \tag{b4} \\
&\quad - (-1)^{|x_2||x_3|+(|x_1|+|x_3|)|f|} (x_1 * x_3) * f(\alpha(x_2)) \tag{a5} \\
&\quad + (-1)^{|x_2||x_3|} f((x_1 * x_3) * \alpha(x_2)) \tag{a6} \\
&\quad - (-1)^{|x_2||x_3|+|x_1||f|} (x_1 * f(x_3)) * \alpha(x_2) \tag{a7} \\
&\quad - (-1)^{|x_2||x_3|} (f(x_1) * x_3) * \alpha(x_2). \tag{a8}
\end{aligned}$$

By (2.5) and (2.6), we have

$$\begin{aligned}
C1 &= (a1) + (b1) - (f(x_1) * x_2) * \alpha(x_3) = 0 \\
C2 &= (a2) + (b2) + f((x_1 * x_2) * \alpha(x_3)) = 0 \\
C3 &= (a3) + (b3) - (-1)^{|x_1||f|} (x_1 * f(x_2)) * \alpha(x_3) = 0 \\
C4 &= (a4) + (b4) - (-1)^{(|x_1|+|x_2|)|f|} (x_1 * x_2) * \alpha(f(x_3)) = 0 \\
C5 &= (a5) - (-1)^{|x_1||f|} (x_1 * f(x_2)) * \alpha(x_3) = 0 \\
C6 &= (a6) - f((x_1 * x_2) * \alpha(x_3)) = 0 \\
C7 &= (a7) + (-1)^{(|x_1|+|x_2|)|f|} (x_1 * x_2) * \alpha(f(x_3)) = 0 \\
C8 &= (a8) + (f(x_1) * x_2) * \alpha(x_3) = 0.
\end{aligned}$$

Thus,

$$\begin{aligned} & \delta_{hom}^2 \delta_{hom}^1 f(x_1, x_2, x_3) \\ &= (a1) + (b1) + (a2) + (b2) + (a3) + (b3) + (a4) + (b4) + (a5) + (a6) + (a7) + (a8) \\ &= C1 + C2 + C3 + C4 + C5 + C6 + C7 + C8 = 0. \end{aligned}$$

The proof is completed. $\square$

For $n = 1, 2$, the map $f \in C_\alpha^n(A, A)$ is called an n -Hom-cocycle if $\delta_{hom}^n f = 0$. We denote by $Z_\alpha^n(A, A)$ the subspace spanned by n -Hom-cocycles and $B_\alpha^n(A, A) = \delta_{hom}^{n-1} C_\alpha^{n-1}(A, A)$. Since $\delta_{hom}^2 \delta_{hom}^1 = 0$, $B_\alpha^2(A, A)$ is a subspace of $Z_\alpha^2(A, A)$. Hence we can define a cohomology space $H_\alpha^2(A, A)$ of $(A, *, \alpha)$ as the factor space $Z_\alpha^2(A, A)/B_\alpha^2(A, A)$.

Let $(A, *, \alpha)$ be a Hom-Novikov superalgebra and $\mathbf{F}[[t]]$ be the ring of formal power series over $\mathbf{F}$. Suppose that $A[[t]]$ is the set of formal power series over A . Then for an $\mathbf{F}$ -bilinear map $f : A \times A \rightarrow A$, it is natural to extend it to be an $\mathbf{F}[[t]]$ -bilinear map $f : A[[t]] \times A[[t]] \rightarrow A[[t]]$ by

$$f \left(\sum_{i \geq 0} x_i t^i, \sum_{j \geq 0} y_j t^j \right) = \sum_{i, j \geq 0} f(x_i, y_j) t^{i+j}.$$

Definition 4.4. Let $(A, *, \alpha)$ be a Hom-Novikov superalgebra over $\mathbf{F}$. A 1-parameter formal deformation of $(A, *, \alpha)$ is a formal power series $g_t : A[[t]] \times A[[t]] \rightarrow A[[t]]$ of the form

$$g_t(x, y) = \sum_{i \geq 0} G_i(x, y) t^i = G_0(x, y) + G_1(x, y) t + G_2(x, y) t^2 + \cdots,$$

where each G_i is an even $\mathbf{F}$ -bilinear map $G_i : A \times A \rightarrow A$ (extended to be $\mathbf{F}[[t]]$ -bilinear) and $G_0(x, y) = x * y$, such that the following identities hold

$$g_t(\alpha(x), \alpha(y)) = \alpha \circ g_t(x, y), \quad (4.2)$$

$$g_t(\alpha(x), g_t(y, z)) - g_t(g_t(x, y), \alpha(z)) = (-1)^{|x||y|} \{g_t(\alpha(y), g_t(x, z)) - g_t(g_t(y, x), \alpha(z))\}, \quad (4.3)$$

$$g_t(g_t(x, y), \alpha(z)) = (-1)^{|y||z|} g_t(g_t(x, z), \alpha(y)). \quad (4.4)$$

Conditions (4.2)-(4.4) are called the deformation equations of a Hom-Novikov superalgebra.

Note that $A[[t]]$ is a module over $\mathbf{F}[[t]]$ and g_t defines the bilinear multiplication on $A[[t]]$ such that $A_t = (A[[t]], g_t, \alpha)$ is a Hom-Novikov superalgebra. Now we investigate the deformation Equations (4.2)-(4.4).

Conditions (4.2)-(4.4) are equivalent to the following equations

$$G_i(\alpha(x), \alpha(y)) = \alpha \circ G_i(x, y), \quad (4.5)$$

respectively, for $i = 0, 1, 2, \dots$. The conditions (4.3) and (4.4) can be expressed as

$$\sum_{i,j \geq 0} \left(G_i(\alpha(x), G_j(y, z)) - G_i(G_j(x, y), \alpha(z)) - (-1)^{|x||y|} G_i(\alpha(y), G_j(x, z)) \right. \\ \left. + (-1)^{|x||y|} G_i(G_j(y, x), \alpha(z)) \right) = 0, \quad (4.6)$$

$$\sum_{i,j \geq 0} \left(G_i(G_j(x, y), \alpha(z)) - (-1)^{|y||z|} G_i(G_j(x, z), \alpha(y)) \right) = 0. \quad (4.7)$$

For $n = 0$, this means $A = A_0$ is a Hom-Novikov superalgebra. For $n = 1$, we obtain three equations for G_1 :

$$G_i(\alpha(x), \alpha(y)) = \alpha \circ G_i(x, y), \quad (4.8)$$

$$G_1(\alpha(x), y * z) - G_1(x * y, \alpha(z)) - (-1)^{|x||y|} G_1(\alpha(y), x * z) + (-1)^{|x||y|} G_1(y * x, \alpha(z)) + \\ \alpha(x) * G_1(y, z) - G_1(x, y) * \alpha(z) - (-1)^{|x||y|} \alpha(y) * G_1(x, z) + (-1)^{|x||y|} G_1(y, x) * \alpha(z) = 0, \quad (4.9)$$

$$(G_1(x * y, \alpha(z)) - (-1)^{|y||z|} G_1(x * z, \alpha(y)) + G_1(x, y) * \alpha(z) - (-1)^{|y||z|} G_1(x, z) * \alpha(y)). \quad (4.10)$$

We call G_1 an infinitesimal deformation.

Example 4.5. A family of Hom-Novikov superalgebras $(A, *_{\xi}, \alpha)$ is an infinitesimal deformation the Hom-Novikov superalgebra defined by Equation (2.9) with a fixed derivation D , where $*_{\xi}$ is defined as

$$x *_{\xi} y = xD(y) + \xi xy, \quad (4.11)$$

for all $x, y \in A$, $\xi \in \mathbf{F}$. For Equation (4.11), we let $G_1(x, y) = xy$, then Equation (4.9) holds since

$$\alpha(x)(yD(z)) - (xD(y))\alpha(z) - (-1)^{|x||y|} \alpha(y)(xD(z)) + (-1)^{|x||y|} (yD(x))\alpha(z) \\ + \alpha(x)D(yz) - (xy)D(\alpha(z)) - (-1)^{|x||y|} \alpha(y)(D(x)z) + (-1)^{|x||y|} (yx)D(\alpha(z)) = 0$$

and Equation (4.10) holds since

$$(xD(y))\alpha(z) - (-1)^{|y||z|} (xD(z))\alpha(y) + (xy)D(\alpha(z)) + (-1)^{|y||z|} (xz)D(\alpha(y)) = 0.$$

Obviously, it is easy to verify that Equation (4.8) holds.

Hence, we have the following result.

Theorem 4.6. Let (A, μ, α) be a Hom-supercommutative algebra and $D : A \rightarrow A$ be an even derivation such that $D\alpha = \alpha D$. Then $(A, *_{\xi}, \alpha)$ is a Hom-Novikov superalgebra, where $*_{\xi}$ is defined as

$$x *_{\xi} y = xD(y) + \xi xy,$$

for all $x, y \in A$, $\xi \in \mathbf{F}$.

For two $\mathbf{F}$ -bilinear maps $f, g : A \times A \rightarrow A$ (extended to be $\mathbf{F}[[t]]$ -bilinear), define a map $f \circ_\alpha g : A[[t]] \times A[[t]] \times A[[t]] \rightarrow A[[t]]$ by

$$f \circ_\alpha g(x, y, z) = f(\alpha(x), g(y, z)) - (-1)^{|x||y|} f(\alpha(y), g(x, z)) + (-1)^{|x||y|} f(g(y, x), \alpha(z)) \\ - (-1)^{|y||z|} f(g(x, z), \alpha(y)).$$

Using Equations (4.6) and (4.7), we have

$$\sum_{i+j=n} G_i \circ_\alpha G_j = 0.$$

For $n = 1$,

$$G_0 \circ_\alpha G_1 + G_1 \circ_\alpha G_0 = 0. \quad (4.12)$$

For $n \geq 2$,

$$-(G_0 \circ_\alpha G_n + G_n \circ_\alpha G_0) = G_1 \circ_\alpha G_{n-1} + G_2 \circ_\alpha G_{n-2} + \cdots + G_{n-1} \circ_\alpha G_1. \quad (4.13)$$

By (4.5) it follows that $G_i \in C_\alpha^2(A, A)$. It can also be verified that $G_i \circ_\alpha G_j \in C_\alpha^3(A, A)$. In general, if $f, g \in C_\alpha^2(A, A)$, then $f \circ_\alpha g \in C_\alpha^3(A, A)$. Note that the definition of coboundary operator, which implies $\delta_{hom}^2 G_n = G_0 \circ_\alpha G_n + G_n \circ_\alpha G_0$, for $n = 0, 1, 2, \dots$. Hence (4.12) and (4.13) can be rewritten as

$$\delta_{hom}^2 G_1 = 0, \\ -\delta_{hom}^2 G_n = G_1 \circ_\alpha G_{n-1} + G_2 \circ_\alpha G_{n-2} + \cdots + G_{n-1} \circ_\alpha G_1.$$

Then G_1 is a 2-Hom-cocycle.

Definition 4.7. Let $(A, *, \alpha)$ be a Hom-Novikov superalgebra. Suppose that $g_t(x, y) = \sum_{i \geq 0} G_i(x, y) t^i$ and $g'_t(x, y) = \sum_{i \geq 0} G'_i(x, y) t^i$ are two 1-parameter formal deformations of $(A, *, \alpha)$. They are called equivalent, denoted by $g_t \sim g'_t$, if there is a formal isomorphism of $\mathbf{F}[[t]]$ -modules

$$\phi_t(x) = \sum_{i \geq 0} \phi_i(x) t^i : (A[[t]], g_t, \alpha) \longrightarrow (A[[t]], g'_t, \alpha),$$

where each $\phi_i : T \rightarrow T$ is an $\mathbf{F}$ -linear map (extended to be $\mathbf{F}[[t]]$ -linear) and $\phi_0 = \text{id}_A$, satisfying

$$\phi_t \circ \alpha = \alpha \circ \phi_t, \\ \phi_t \circ g_t(x, y) = g'_t(\phi_t(x), \phi_t(y)).$$

When $G_1 = G_2 = \cdots = 0$, $g_t = G_0$ is said to be the null deformation. A 1-parameter formal deformation g_t is called trivial if $g_t \sim G_0$. A Hom-Novikov superalgebra $(T, [\cdot, \cdot], \alpha)$ is called analytically rigid, if every 1-parameter formal deformation g_t is trivial.

Theorem 4.8. Let $g_t(x, y, z) = \sum_{i \geq 0} G_i(x, y, z)t^i$ and $g'_t(x, y, z) = \sum_{i \geq 0} G'_i(x, y, z)t^i$ be equivalent 1-parameter formal deformations of $(A, *, \alpha)$. Then G_1 and G'_1 belong to the same cohomology class in $H^2_\alpha(A, A)$.

Proof. Suppose that $\phi_t(x) = \sum_{i \geq 0} \phi_i(x)t^i$ is the formal $\mathbf{F}[[t]]$ -module isomorphism such that $\phi_t \circ \alpha = \alpha \circ \phi_t$ and

$$\sum_{i \geq 0} \phi_i \left(\sum_{j \geq 0} G_j(x, y)t^j \right) t^i = \sum_{i \geq 0} G'_i \left(\sum_{k \geq 0} \phi_k(x)t^k, \sum_{l \geq 0} \phi_l(y)t^l \right) t^i.$$

It follows that

$$\sum_{i+j=n} \phi_i(G_j(x, y))t^{i+j} = \sum_{i+k+l=n} G'_i(\phi_k(x), \phi_l(y))t^{i+k+l}.$$

In particular,

$$\sum_{i+j=1} \phi_i(G_j(x, y)) = \sum_{i+k+l=1} G'_i(\phi_k(x), \phi_l(y)),$$

that is,

$$G_1(x, y) + \phi_1(x * y) = \phi_1(x) * y + x * \phi_1(y) + G'_1(x, y).$$

Then $G_1 - G'_1 = \delta_{hom}^1 \phi_1 \in B_\alpha^2(A, A)$. □

Theorem 4.9. Suppose that $(A, *, \alpha)$ is a Hom-Novikov superalgebra such that $H_\alpha^2(A, A) = 0$. Then $(A, *, \alpha)$ is analytically rigid.

Proof. Let g_t be a 1-parameter formal deformation of $(A, *, \alpha)$. Suppose that $g_t = G_0 + \sum_{i \geq n} G_i t^i$. Then

$$\delta_{hom}^2 G_n = G_1 \circ_\alpha G_{n-1} + G_2 \circ_\alpha G_{n-2} + \cdots + G_{n-1} \circ_\alpha G_1 = 0,$$

that is, $G_n \in Z_\alpha^2(A, A) = B_\alpha^2(A, A)$. It follows that there exists $f_n \in C_\alpha^1(A, A)$ such that $G_n = \delta_{hom}^1 f_n$.

Let $\phi_t = \text{id}_A - f_n t^n : (A[[t]], g'_t, \alpha) \longrightarrow (A[[t]], g_t, \alpha)$. Note that

$$\phi_t \circ \sum_{i \geq 0} f_n^i t^{in} = \sum_{i \geq 0} f_n^i t^{in} \circ \phi_t = \text{id}_{A[[t]]}.$$

Then ϕ_t is a linear isomorphism. Moreover, $\phi_t \circ \alpha = \alpha \circ \phi_t$.

Now consider $g'_t(x, y) = \phi_t^{-1} g_t(\phi_t(x), \phi_t(y))$. It is straightforward to prove that g'_t is a 1-parameter formal deformation of $(A, *, \alpha)$ and $g_t \sim g'_t$. Suppose that $g'_t = \sum_{i \geq 0} G'_i t^i$. Then

$$(\text{id}_T - f_n t^n) \left(\sum_{i \geq 0} G'_i(x, y)t^i \right) = \left(G_0 + \sum_{i \geq n} G_i t^i \right) (x - f_n(x)t^n, y - f_n(y)t^n),$$

i.e.,

$$\begin{aligned}
& \sum_{i \geq 0} G'_i(x, y) t^i - \sum_{i \geq 0} f_n \circ G'_i(x, y) t^{i+n} \\
&= x * y - (f_n(x) * y + x * f_n(y)) t^n + f_n(x) * f_n(y) t^{2n} \\
&+ \sum_{i \geq n} G_i(x, y) t^i - \sum_{i \geq n} (G_i(f_n(x), y) + G_i(x, f_n(y))) t^{i+n} + \sum_{i \geq n} G_i(f_n(x), f_n(y)) t^{i+2n}.
\end{aligned}$$

Then we have $G'_1 = \cdots = G'_{n-1} = 0$ and

$$G'_n(x, y) - f_n(x * y) = -(f_n(x) * y + x * f_n(y)) + G_n(x, y).$$

Hence $G'_n = G_n - \delta_{hom}^1 f_n = 0$ and $g'_t = G_0 + \sum_{i \geq n+1} G'_i t^i$. By induction, this procedure ends with $g_t \sim G_0$, i.e., $(A, *, \alpha)$ is analytically rigid. $\square$

References

- [1] F. Ammar, A. Makhlouf. Hom-Lie superalgebras and Hom-Lie admissible superalgebras. J. Algebra 324 (2010), 1513–1528.
- [2] C. Bai, D. Meng. A Lie algebraic approach to Novikov algebras. J. Geom. Phys. 45 (2003), 218–230.
- [3] C. Bai, D. Meng. On the Novikov algebra structures adapted to the automorphism structure of a Lie group. J. Geom. Phys. 45 (2003), 105–115.
- [4] C. Bai, D. Meng. The realization of non-transitive Novikov algebras. J. Phys. A 34 (2001), 6435–6442.
- [5] L. Chen, Y. Niu, D. Meng. Two kinds of Novikov algebras and their realizations. J. Pure Appl. Algebra 212 (2008), 902–909.
- [6] L. Chen. A kind of infinite-dimensional Novikov algebras and its realizations. Abstr. Appl. Anal. (2013), 270937, 5pp.
- [7] I. Gel'fand, L. Diki. Asymptotic behaviour of the resolvent of Strum-Liouville equations and the algebra of the Korteweg-deVries equations. Russian Math. Surveys. 30 (1975), 77–113.
- [8] I. Gel'fand, L. Diki. A Lie algebra structure in a formal variational calculation. Funct. Anal. Appl. 10 (1976), 16–22.
- [9] I. Gel'fand, I. Dorfman. Hamiltonian operators and algebraic structures related to them. Funktsional. Anal. i Prilozhen. 13 (1979), 13–30.
- [10] M. Gerstenhaber. On the deformation of rings and algebras. Ann. of Math. (2) 79 (1964), 59–103.

- [11] M. Gerstenhaber. On the deformation of rings and algebras. II. *Ann. of Math.* 84 (1966), 1–19.
- [12] M. Gerstenhaber. On the deformation of rings and algebras. III. *Ann. of Math.* (2) 88 (1968), 1–34.
- [13] M. Gerstenhaber. On the deformation of rings and algebras. IV. *Ann. of Math.* (2) 99 (1974), 257–276.
- [14] Y. Kang, Z. Chen. Novikov superalgebras in low dimensions. *J. Nonlinear math. Phys.* 16 (2009), 251–257.
- [15] F. Kubo, Y. Taniguchi. A controlling cohomology of the deformation theory of Lie triple systems. *J. Algebra* 278 (2004), 242–250.
- [16] Y. Liu, L. Chen, Y. Ma. Hom-Nijenhuis operators and T^* -extensions of hom-Lie superalgebras. *Linear Algebra Appl.* 439 (2013), 2131–2144.
- [17] Y. Ma, L. Chen, J. Lin. One-parameter formal deformations of Hom-Lie Yamaguti algebras. *J. Math. Phys.* in press.
- [18] Y. Ma, L. Chen, J. Lin. Cohomology and 1-parameter formal deformations of Hom-Lie triple systems. *arXiv:13093347*.
- [19] J. Ni, Z. Chen. Novikov super-algebras with associative nondegenerate super-symmetric bilinear forms. *J. Nonlinear Math. Phys.* 17 (2010), 159–166.
- [20] J. Osborn. Infinite dimensional Novikov algebras of characteristic 0. *J. Algebra* 16 (1994), 146–167.
- [21] J. Osborn. simple Novikov algebras with an idempotent. *Commun. Algebra* 20 (1992), 2729–2753.
- [22] X. Xu. Variational calculus of supervariables and related algebraic structures. *J. Algebra* 223 (2000), 396–437.
- [23] X. Xu. Hamiltonian operators and associative algebras with a derivation. *Lett. Math. Phys.* 33 (1995), 1–6.
- [24] X. Xu. On simple Novikov algebra and their irreducible modules. *J. Algebra* 185 (1996), 905–934.
- [25] X. Xu. Novikov-Poisson algebras. *J. Algebra* 190 (1997), 253–279.
- [26] D. Yau. Hom-Novikov algebras. *J. Phys. A: Math. Theor.* 44 (2011), 085202, 20pp.
- [27] L. Yuan, H. You. Hom-Novikov algebras and Hom-Novikov-Poisson algebras. *arXiv:1204.6373*.

- [28] R. Zhang, D. Hou, C. Bai. A Hom-version of the affinizations of Balinskii-Novikov and Novikov superalgebras. *J. Math. Phys.* 52 (2011), 023505, 19 pp.
- [29] E. Zel'manov. On a class of local translation invariant Lie algebras. *Soviet Math. Dokl.* 35 (1987), 216–218.
- [30] F. Zhu, Z. Chen. Novikov superalgebras with $A_0 = A_1A_1$. *Czechoslovak Mathematical Journal* 60 (2010), 903–907.